

# Global Fits with Sterile Neutrinos

Joachim Kopp (University of Mainz)

CrossTalk Workshop “The Fate of Sterile Neutrinos”, January 18, 2017



# Outline

- 1 Sterile Neutrinos
- 2 Oscillation Anomalies: A Global Fit
  - $\nu_e$  Appearance
  - $\nu_e$  Disappearance
  - $\nu_\mu$  Disappearance
  - Sterile Neutrino Oscillations: The Global Picture
- 3 Sterile Neutrinos in Cosmology: Robustness of Constraints
- 4 Light Sterile Neutrinos and Dark Matter Searches
  - Sterile Neutrinos and Direct Dark Matter Searches
  - Sterile Neutrinos and Indirect Dark Matter Searches
- 5 Summary



# Sterile Neutrinos

# Sterile neutrinos

## Definition

Sterile neutrino = SM singlet fermion

- Very generic extension of the SM
  - ▶ can be leftovers of **extended gauge multiplets** (e.g. GUT multiplets)
- Very useful in phenomenology:
  - ▶ Can explain **smallness of neutrino mass** (seesaw mechanism,  $m \sim \text{TeV} \dots M_{\text{Pl}}$ )
  - ▶ Can explain **baryon asymmetry of the Universe** (leptogenesis,  $m \gg 100 \text{ GeV}$ )
  - ▶ Can explain **dark matter** ( $m \sim \text{keV}$ )
  - ▶ Can explain various **neutrino oscillation anomalies** ( $m \sim \text{eV}$ )






## Oscillation Anomalies: A Global Fit


# $\bar{\nu}_e$ appearance in the 3+1 scenario and beyond

Motivated by LSND and MiniBooNE: excess of  $\bar{\nu}_e$  events in  $\bar{\nu}_\mu$  beam.



	$\chi^2_{3+1}/\text{dof}$	$\chi^2_{3+2}/\text{dof}$	$\chi^2_{1+3+1}/\text{dof}$
LSND	11.0/11	8.6/11	7.5/11
MiniB $\nu$	19.3/11	10.6/11	9.1/11
MiniB $\bar{\nu}$	10.7/11	9.6/11	12.7/11
E776	32.4/24	29.2/24	31.3/24
KARMEN	9.8/9	8.6/9	9.0/9
NOMAD	0.0/1	0.0/1	0.0/1
ICARUS	2.0/1	2.3/1	1.5/1
<b>Combined</b>	<b>87.9/66</b>	<b>72.7/63</b>	<b>74.6/63</b>


- Global fit to all appearance data is **consistent**
- **Background oscillations** important in MiniBooNE and E776
- Significant improvement in **3 + 2** and **1 + 3 + 1**



Dentler JK Machado Maltoni Schwetz, in prep.  
see also fits by Giunti Laveder et al.  
Collin Argüelles Conrad Shaevitz


# $\bar{\nu}_e$ appearance in the 3+1 scenario and beyond

Motivated by LSND and MiniBooNE: excess of  $\bar{\nu}_e$  events in  $\bar{\nu}_\mu$  beam.



	$\chi^2_{3+1}/\text{dof}$	$\chi^2_{3+2}/\text{dof}$	$\chi^2_{1+3+1}/\text{dof}$
LSND	11.0/11	8.6/11	7.5/11
MiniB $\nu$	19.3/11	10.6/11	9.1/11
MiniB $\bar{\nu}$	10.7/11	9.6/11	12.7/11
E776	32.4/24	29.2/24	31.3/24
KARMEN	9.8/9	8.6/9	9.0/9
NOMAD	0.0/1	0.0/1	0.0/1
ICARUS	2.0/1	2.3/1	1.5/1
<b>Combined</b>	<b>87.9/66</b>	<b>72.7/63</b>	<b>74.6/63</b>


- Global fit to all appearance data is **consistent**
- **Background oscillations** important in MiniBooNE and E776
- Significant improvement in **3 + 2** and **1 + 3 + 1**



Dentler JK Machado Maltoni Schwetz, in prep.  
see also fits by Giunti Laveder et al.  
Collin Argüelles Conrad Shaevitz


# $\bar{\nu}_e$ appearance in the 3+1 scenario and beyond

Motivated by LSND and MiniBooNE: excess of  $\bar{\nu}_e$  events in  $\bar{\nu}_\mu$  beam.



	$\chi^2_{3+1}/\text{dof}$	$\chi^2_{3+2}/\text{dof}$	$\chi^2_{1+3+1}/\text{dof}$
LSND	11.0/11	8.6/11	7.5/11
MiniB $\nu$	19.3/11	10.6/11	9.1/11
MiniB $\bar{\nu}$	10.7/11	9.6/11	12.7/11
E776	32.4/24	29.2/24	31.3/24
KARMEN	9.8/9	8.6/9	9.0/9
NOMAD	0.0/1	0.0/1	0.0/1
ICARUS	2.0/1	2.3/1	1.5/1
<b>Combined</b>	<b>87.9/66</b>	<b>72.7/63</b>	<b>74.6/63</b>

- Global fit to all appearance data is consistent
- Background oscillations important in MiniBooNE and E776
- Significant improvement in 3 + 2 and 1 + 3 + 1



Dentler JK Machado Maltoni Schwetz, in prep.  
see also fits by Giunti Laveder et al.  
Collin Argüelles Conrad Shaevitz




# $\bar{\nu}_e$ disappearance: reactor and gallium experiments

## The reactor anomaly

- **Reevaluation** of reactor  $\bar{\nu}_e$  flux is  $\sim 3.5\%$  **above** previous prediction  
→ **systematic uncertainties** or **new physics**?

Mueller et al. 1101.2663, Huber 1106.0687, Hayes et al. 1309.4146 & talk by G. Garvey



# $\bar{\nu}_e$ disappearance: reactor and gallium experiments

## The reactor anomaly


- **Reevaluation** of reactor  $\bar{\nu}_e$  flux is  $\sim 3.5\%$  **above** previous prediction  
→ **systematic uncertainties** or **new physics**?

Mueller et al. 1101.2663, Huber 1106.0687, Hayes et al. 1309.4146 & talk by [G. Garvey](#)



## The gallium anomaly

- Experiments with **intense radioactive  $\nu_e$  sources** ( $^{51}\text{Cr}$  and  $^{37}\text{Ar}$ )
- Neutrino detection via  
 $^{71}\text{Ga} + \nu_e \rightarrow ^{71}\text{Ge} + e^-$
- **Observation: Neutrino deficit** ( $\sim 3\sigma$ )

Giunti Laveder 1006.3244





# $\bar{\nu}_e$ disappearance in the 3+1 scenario



	$\sin^2 2\theta_{14}$	$\Delta m_{41}^2 [\text{eV}^2]$	$\chi_{\min}^2/\text{dof}$ (GOF)	$\Delta\chi_{\text{no osc}}^2/\text{dof}$ (CL)
SBL rates only	0.13	0.44	11.5/17 (83%)	11.4/2 (99.7%)
SBL incl. Bugey3 spect.	0.10	1.75	58.3/74 (91%)	9.0/2 (98.9%)
SBL + Gallium	0.11	1.80	64.0/78 (87%)	14.0/2 (99.9%)
global $\nu_e$ disapp.	0.09	1.78	403.3/427 (79%)	12.6/2 (99.8%)

Dentler JK Machado Maltoni Schwetz, in prep.

# $\bar{\nu}_e$ disappearance in the 3+1 scenario



	$\sin^2 2\theta_{14}$	$\Delta m_{41}^2 [\text{eV}^2]$	$\chi_{\min}^2/\text{dof}$ (GOF)	$\Delta\chi_{\text{no osc}}^2/\text{dof}$ (CL)
SBL rates only	0.13	0.44	11.5/17 (83%)	11.4/2 (99.7%)
SBL incl. Bugey3 spect.	0.10	1.75	58.3/74 (91%)	9.0/2 (98.9%)
SBL + Gallium	0.11	1.80	64.0/78 (87%)	14.0/2 (99.9%)
global $\nu_e$ disapp.	0.09	1.78	403.3/427 (79%)	12.6/2 (99.8%)

Dentler JK Machado Maltoni Schwetz, in prep.

# Relation between appearance and disappearance

We find:  $\bar{\nu}_e$  disappearance experiments consistent among themselves,  
 $\bar{\nu}_e$  appearance experiments consistent among themselves.

But:

## 3 + 1 neutrinos

At  $L \gg 4\pi E/\Delta m_{41}^2$ , but  $L \ll 4\pi E/\Delta m_{31}^2$

$$P_{ee} = 1 - 2|U_{e4}|^2(1 - |U_{e4}|^2)$$

$$P_{\mu\mu} = 1 - 2|U_{\mu4}|^2(1 - |U_{\mu4}|^2)$$

$$P_{e\mu} = 2|U_{e4}|^2|U_{\mu4}|^2$$

It follows


$$2P_{e\mu} \simeq (1 - P_{ee})(1 - P_{\mu\mu})$$

In the 3 + 1 case, at **large enough baseline**, there is a **one-to-one relation** between the **appearance and disappearance probabilities**.

Combining different oscillation channels  
provides the **strongest, most robust**  
constraints on sterile neutrinos


# $\bar{\nu}_\mu$ disappearance in the 3+1 scenario (as of 2013)

- Parameter regions favored by **tentative hints** are in **tension with null results** from  $\bar{\nu}_\mu$  disappearance searches



JK Machado Maltoni Schwetz, 1303.3011

# IceCube sterile neutrino search




IceCube arXiv:1605.01990


- New technique to search for  $\nu_s$
- MSW resonance between  $\nu_a$  and  $\nu_s$ 
  - ▶ at  $E \sim \text{TeV}$  for  $\Delta m_{41}^2 \sim \text{eV}$
- Unique spectral feature
- But only in  $\bar{\nu}$



# IceCube sterile neutrino search




IceCube arXiv:1605.01990




Dentler JK Machado Maltoni Martinez Schwetz,  
in preparation

# $\bar{\nu}_\mu$ disappearance in the 3+1 scenario



Dentler JK Machado Maltoni Martinez Schwetz, in preparation  
Plot courtesy of Mona Dentler



# $\bar{\nu}_\mu$ disappearance in the 3+1 scenario



Dentler JK Machado Maltoni Martinez Schwetz, in preparation  
Plot courtesy of Mona Dentler

# $\bar{\nu}_\mu$ disappearance in the 3+1 scenario

- Constraints on  $|U_{\tau 4}| \sim \sin \theta_{34}$  from NC events and matter effects (2013)




JK Machado Maltoni Schwetz, arXiv:1303.3011

# The global oscillation fit (as of 2013)

JK Machado Maltoni Schwetz, arXiv:1303.3011

3 + 1 Severe **tension** between appearance and disappearance and between exp's with and without a signal




	$\chi^2_{\min}/\text{dof}$	GOF
3+1	712/(689 - 9)	19%

# The global oscillation fit (as of 2013)

JK Machado Maltoni Schwetz, arXiv:1303.3011

3 + 1 Severe **tension** between appearance and disappearance and between exp's with and without a signal



Parameter goodness of fit (PG) test:

Compares  $\chi^2_{\min}$  from global and separate fits to test **compatibility of 2 data sets**


Maltoni Schwetz hep-ph/0304176

	$\chi^2_{\min}/\text{dof}$	GOF	$\chi^2_{\text{PG}}/\text{dof}$	PG
3+1	712/(689 - 9)	19%	18.0/2	$1.2 \times 10^{-4}$

# The global oscillation fit (as of 2013)

JK Machado Maltoni Schwetz, arXiv:1303.3011

3 + 1 Severe **tension** between appearance and disappearance and between exp's with and without a signal



Parameter goodness of fit (PG) test:

Compares  $\chi^2_{\min}$  from global and separate fits to test **compatibility of 2 data sets**


Maltoni Schwetz hep-ph/0304176

	$\chi^2_{\min}/\text{dof}$	GOF	$\chi^2_{\text{PG}}/\text{dof}$	PG
3+1 (+IC)	712/(689 - 9)	19%	18.0/2	$0.4 \times 10^{-4}$

# The global oscillation fit (as of 2013)

JK Machado Maltoni Schwetz, arXiv:1303.3011

- 3 + 1 Severe **tension** between appearance and disappearance and between exp's with and without a signal
- 3 + 2 **Tension remains** for **two sterile neutrinos**



Parameter goodness of fit (PG) test:

Compares  $\chi^2_{\min}$  from global and separate fits to test **compatibility of 2 data sets**

Maltoni Schwetz hep-ph/0304176


	$\chi^2_{\min}/\text{dof}$	GOF	$\chi^2_{\text{PG}}/\text{dof}$	PG
3+1 (+IC)	712/(689 - 9)	19%	18.0/2	$0.4 \times 10^{-4}$
3+2	701/(689 - 14)	23%	25.8/4	$3.4 \times 10^{-5}$
1+3+1	694/(689 - 14)	30%	16.8/4	$2.1 \times 10^{-3}$



# The global oscillation fit (as of 2013)

JK Machado Maltoni Schwetz, arXiv:1303.3011

- 3 + 1 Severe **tension** between appearance and **disappearance** and between exp's with and without a signal
- 3 + 2 **Tension remains** for **two sterile neutrinos**
- 3 + 3 No significant improvement expected



Parameter goodness of fit (PG) test:

Compares  $\chi^2_{\min}$  from global and separate fits to test **compatibility of 2 data sets**

Maltoni Schwetz hep-ph/0304176

	$\chi^2_{\min}/\text{dof}$	GOF	$\chi^2_{\text{PG}}/\text{dof}$	PG
3+1 (+IC)	712/(689 - 9)	19%	18.0/2	$0.4 \times 10^{-4}$
3+2	701/(689 - 14)	23%	25.8/4	$3.4 \times 10^{-5}$
1+3+1	694/(689 - 14)	30%	16.8/4	$2.1 \times 10^{-3}$

# The MIT/Columbia fit

- $(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$  appearance data:

- ▶ LSND
- ▶ MiniBooNE
- ▶ KARMEN
- ▶ NOMAD

- $(\bar{\nu}_\mu)$  disappearance data:

- ▶ MiniBooNE
- ▶ Minos CC  $\nu_\mu$
- ▶ CDHS
- ▶ CCFR
- ▶ Atmospheric neutrinos
- ▶ IceCube



- $(\bar{\nu}_e)$  disappearance data:

- ▶ Short baseline reactor experiments
- ▶ Gallium experiments
- ▶  $\nu_e$ - $^{12}\text{C}$  CC scattering in KARMEN, LSND

Collin Argüelles Conrad Shaevitz, arXiv:1607.00011, 1602.00671v1  
 Conrad Ignarra Karagiorgi Shaevitz Spitz, arXiv:1207.4765


Authors **question applicability** of parameter goodness of fit test

$$|U| = \begin{bmatrix} 0.79 \rightarrow 0.83 & 0.53 \rightarrow 0.57 & 0.14 \rightarrow 0.15 & 0.13 \rightarrow 0.20 \\ 0.25 \rightarrow 0.50 & 0.46 \rightarrow 0.66 & 0.64 \rightarrow 0.77 & 0.09 \rightarrow 0.15 \\ 0.26 \rightarrow 0.54 & 0.48 \rightarrow 0.69 & 0.56 \rightarrow 0.75 & 0.0 \rightarrow 0.7 \\ \dots & \dots & \dots & \dots \end{bmatrix}$$



# The $GL^4$ fit

- $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  appearance data:
  - ▶ LSND
  - ▶ MiniBooNE
  - ▶ E776
  - ▶ KARMEN
  - ▶ NOMAD
  - ▶ ICARUS
  - ▶ OPERA
- $\bar{\nu}_\mu$  disappearance data:
  - ▶ MiniBooNE/SciBooNE
  - ▶ Minos NC+CC  $\nu_\mu$
  - ▶ CDHS
  - ▶ CCFR
  - ▶ Atmospheric neutrinos
- $\bar{\nu}_e$  disappearance data:
  - ▶ Reactor experiments
  - ▶ Gallium experiments
  - ▶ Solar neutrinos
  - ▶  $\nu_e$ - $^{12}\text{C}$  scattering in KARMEN, LSND



Giunti, Neutrino 2016, arXiv:1609.04688  
Giunti Laveder Li Long arXiv:1308.5288  
Giunti Laveder Li Liu Long arXiv:1210.5715  
Giunti Laveder arXiv:1111.1069

Conclusion

Tension confirmed

## Conclusion from oscillation data:

### severe tension


#### My personal point of view

Anomalies are definitely interesting enough to be checked

- If **mundane explanations** are found, we will learn **important lessons for future neutrino experiments**
- If **sterile neutrinos** exist, this would be the **discovery of the decade**.

#### Questions for the future

- Is one (or all) of the **positive results** not due to sterile neutrinos?
- Are some of the **exclusion limits** too strong?  
(removing one from the fit is not enough.)
- Does **new physics beyond the simple  $3 + X$  scenarios** help?



# Sterile Neutrinos in Cosmology: Robustness of Constraints

# Sterile neutrinos in cosmology

Models with  $\mathcal{O}(\text{eV})$  sterile neutrino(s) constrained by cosmology:

Sum of neutrino masses

$$\sum m_\nu \lesssim 0.23 \text{ eV}$$

# of relativistic species

$$N_\nu = 4 \text{ mildly disfavored}$$

Ade et al. (Planck), arXiv:1303.5076

Gonzalez-Garcia Maltoni Salvado, arXiv:1006.3795

Hamann Hannestad Raffelt Tamborra Wong, arXiv:1006.5276

# Sterile neutrinos in cosmology

Models with  $\mathcal{O}(\text{eV})$  sterile neutrino(s) constrained by cosmology:

Sum of neutrino masses

$$\sum m_\nu \lesssim 0.23 \text{ eV}$$

# of relativistic species

$$N_\nu = 4 \text{ mildly disfavored}$$

Ade et al. (Planck), arXiv:1303.5076

Gonzalez-Garcia Maltoni Salvado, arXiv:1006.3795

Hamann Hannestad Raffelt Tamborra Wong, arXiv:1006.5276

Question:

What does it take to **evade** these constraints?

# Are light sterile neutrinos ruled out by cosmology?

$\nu_s$  production in the early Universe through  $\nu_{e,\mu,\tau} \rightarrow \nu_s$  oscillations at  $T \gtrsim \text{MeV}$

Dodelson Widrow 1994



## Making sterile neutrinos fully consistent with cosmology

- $> 1$  new relativistic degrees of freedom +  $w < -1$  +  $\mu_\nu \neq 0$   
Hamann Hannestad Raffelt Wong, arXiv:1108.4136
- **Entropy production** after neutrino decoupling (e.g. due to late decay of heavy sterile neutrinos or other particles)  $\rightarrow$  neutrinos diluted  
Fuller Kishimoto Kusenko 1110.6479, Ho Scherrer 1212.1689
- Very low reheating temperature  
Gelmini Palomares-Ruiz Pascoli, astro-ph/0403323
- Large **lepton asymmetry** ( $\gtrsim 0.01$ )  $\rightarrow \nu_s$  production MSW-suppressed  
Foot Volkas hep-ph/9508275, Chu Cirelli astro-ph/0608206, Saviano et al. arXiv:1302.1200
- Couplings to a **Majoron field**  $\rightarrow$  suppressed production  
Bento Berezhiani, hep-ph/0108064
- **New gauge interaction** in the  $\nu_s$  sector  
 $\rightarrow \nu_s$  production suppressed by thermal potential  
Hannestad et al. 1310.5926  
Dasgupta JK 1310.6337



# Suppressed $\nu_s$ production from thermal MSW effect

- $\nu_s$  production in the early Universe through  $\nu_{e,\mu,\tau} \rightarrow \nu_s$  oscillations  
Dodelson Widrow 1994
- Assume  $\nu_s$  charged under a **hidden sector gauge group**  $U(1)_s$
- Neutrino **self energy**:





- Tadpole digram**: conventional MSW potential:  $V \propto n_f - n_{\bar{f}}$   
**Bubble digram**: thermal contribution  $V \propto T^\alpha$
- At high  $T$ : strong MSW-type potential **even without lepton asymmetry**

$$\sin 2\theta_{\text{eff}} = \frac{\sin 2\theta}{\sqrt{\sin^2 2\theta + \left(\cos 2\theta - \frac{2EV}{\Delta m^2}\right)^2}}$$

- $\nu_s$  production through oscillations **suppressed** in the early Universe  
→ **no cosmological constraints**  
Hannestad Hansen Tram arXiv:1310.5926  
Dasgupta JK arXiv:1310.6337

# Suppression of $\nu_s$ production by thermal MSW effect



- If  $V_{\text{eff}} \gg \Delta m^2 / (2E)$  until decoupling:  
 $\Rightarrow$  **sterile neutrino production suppressed**, no cosmological constraints

Hannestad Hansen *Tram* arXiv:1310.5926

Dasgupta JK arXiv:1310.6337

- Details still under investigation

Chu Dasgupta JK arXiv:1505.02795

Cherry Friedland Shoemaker arXiv:1605.06506

Dentler JK Saviano, in preparation

The Google logo is centered at the top of the slide. It consists of the word "Google" in its characteristic multi-colored font: 'G' is blue, 'o' is red, 'o' is yellow, 'g' is green, 'l' is blue, and 'e' is red.A search input field with a light blue border and a thin shadow. The text "Dark Matter" is entered into the field, followed by a vertical cursor line.A rectangular button with a light gray background and a thin border. The text "Google Search" is centered on the button.A rectangular button with a light gray background and a thin border. The text "I'm Feeling Lucky" is centered on the button.



# Light Sterile Neutrinos and Dark Matter Searches

# Neutrinos and direct dark matter detection

Solar neutrinos are a well-known **background** to future direct DM searches:

see e.g. Gütlein et al. arXiv:1003.5530

$$\frac{d\sigma_{\text{SM}}(\nu N \rightarrow \nu N)}{dE_r} = \frac{G_F^2 m_N F^2(E_r)}{2\pi E_\nu^2} \left[ A^2 E_\nu^2 + 2AZ(2E_\nu^2(s_W^2 - 1) - E_r m_N s_W^2) + 4Z^2(E_\nu^2 + s_W^4(2E_\nu^2 + E_r^2 - E_r(2E_\nu + m_N)) + s_W^2(E_r m_N - 2E_\nu^2)) \right],$$



# Neutrinos and direct dark matter detection

Solar neutrinos are a well-known **background** to future direct DM searches:

see e.g. Gütlein et al. arXiv:1003.5530

$$\frac{d\sigma_{\text{SM}}(\nu N \rightarrow \nu N)}{dE_r} = \frac{G_F^2 m_N F^2(E_r)}{2\pi E_\nu^2} \left[ A^2 E_\nu^2 + 2AZ(2E_\nu^2(s_w^2 - 1) - E_r m_N s_w^2) \right. \\ \left. + 4Z^2(E_\nu^2 + s_w^4(2E_\nu^2 + E_r^2 - E_r(2E_\nu + m_N)) + s_w^2(E_r m_N - 2E_\nu^2)) \right],$$


SM signal will only become sizeable in **multi-ton detectors**

**But:** New physics can **enhance** the rate

Examples:


- **Neutrino magnetic moments**
- **Sterile neutrinos** +  $\lesssim$  **GeV scale hidden sector gauge force**

# Low-energy scattering of neutrinos beyond the SM



A:  $\nu$  magnetic moment  
 B, C, D: kinetically mixed  $A'$  + sterile  $\nu_s$

- Enhanced scattering at low  $E_r$  for light  $A'$
- Negligible compared to SM scattering ( $\sim g^4 m_T/M_W^4$ ) at energies probed in dedicated neutrino experiments



A:  $\nu$  magnetic moment  
 B:  $U(1)_{B-L}$  boson  
 C: kinetically mixed  $U(1)'$  + sterile  $\nu$   
 D:  $U(1)_B$  + sterile  $\nu$  charged under  $U(1)_B$   
 [Pospelov 1103.3261, Pospelov Pradler 1203.0545]


[Harnik JK Machado arXiv:1202:6073]

# Sterile neutrinos and DM annihilation in the Sun

- Neutrino telescope limits on **neutrinos from dark matter annihilation in the Sun** depend crucially on oscillation physics.
- If **sterile neutrinos** exist, new **MSW resonances** can lead to **strong conversion of active neutrinos into sterile neutrinos** in the Sun.

Esmaili Peres, arXiv:1202.2869  
Argüelles JK, arXiv:1202.3431

# Oscillation probabilities for a 3+3 toy scenario




Thick red lines = active-sterile oscillations

plots from Argüelles JK, arXiv:1202.3431  
see also Esmaili Peres, arXiv:1202.2869




# Impact on IceCube limits



plot from Carlos Argüelles JK, arXiv:1202.3431  
see also Esmaili Peres, arXiv:1202.2869

# Flavor ratios in the presence of sterile neutrinos



- Allowed ranges largely unchanged when global constraints on  $\nu_s$  mixing are taken into account
- Interesting possibility: Primary flux of  $\nu_s$ , for instance from dark matter decay
- For the future: flavor ratios binned in energy?

Brdar JK Wang, in preparation



# Summary

# Summary

- Sterile neutrinos are **theoretically well motivated** and **phenomenologically useful**
- **Tension** between **appearance** and **disappearance** searches
- **IceCube** increases tension significantly
  - ▶ Completely new method
  - ▶ Systematics very different from other  $\nu_s$  searches
  - ▶ Theory community very excited about update with more data
- Cosmology **disfavors**  $N_\nu \geq 4$ , especially for  $m_{\nu_s} \gtrsim 0.23$  eV.  
(if BICEP-2 is included, this conclusion would change)
- Cosmology is not a show-stopper
  - ▶ Example: **hidden sector gauge force**
- Sterile neutrinos and **dark matter searches**
  - ▶ **Direct searches**: **non-standard neutrino signals** in DM detectors
  - ▶ **Indirect searches**: limits on **DM annihilation in the Sun** modified by active–sterile oscillations



Thank you!

# Data sets included in our fit

## $\bar{\nu}_e$ disappearance

- SBL reactor experiments
- LBL reactor experiments
- KamLAND
- Radioactive source (Ga) experiments
- Solar neutrinos
- Atmospheric neutrinos
- $\nu_e$ - $^{12}\text{C}$  scattering in KARMEN, LSND

## $\bar{\nu}_e$ appearance

- LSND
- MiniBooNE
- KARMEN
- NOMAD
- ICARUS
- E776

## $\bar{\nu}_\mu$ disappearance

- Atmospheric neutrinos (includes *either*  $\bar{\nu}_e$  disapp. *or* full matter effects)
- MiniBooNE (includes oscillations of backgrounds)
- MINOS CC+NC (full  $n$ -flavour oscillations in matter)
- CDHS

# Relation between appearance and disappearance

## 3 + 2 neutrinos

At  $L \gg 4\pi E/\Delta m_{41}^2$ , but  $L \ll 4\pi E/\Delta m_{31}^2$

$$P_{ee} = 1 - 2 \left[ |U_{e4}|^2(1 - |U_{e4}|^2) + |U_{e5}|^2(1 - |U_{e5}|^2) - |U_{e4}|^2|U_{e5}|^2 \right]$$

$$P_{\mu\mu} = 1 - 2 \left[ |U_{\mu4}|^2(1 - |U_{\mu4}|^2) + |U_{\mu5}|^2(1 - |U_{\mu5}|^2) - |U_{\mu4}|^2|U_{\mu5}|^2 \right]$$

$$P_{e\mu} = 2 \left[ |U_{e4}|^2|U_{\mu4}|^2 + |U_{e5}|^2|U_{\mu5}|^2 + \text{Re}(U_{e4}^* U_{\mu4} U_{e5} U_{\mu5}^*) \right]$$

# Relation between appearance and disappearance

## 3 + 2 neutrinos

At  $L \gg 4\pi E/\Delta m_{41}^2$ , but  $L \ll 4\pi E/\Delta m_{31}^2$

$$P_{ee} = 1 - 2 \left[ |U_{e4}|^2(1 - |U_{e4}|^2) + |U_{e5}|^2(1 - |U_{e5}|^2) - |U_{e4}|^2|U_{e5}|^2 \right]$$

$$P_{\mu\mu} = 1 - 2 \left[ |U_{\mu4}|^2(1 - |U_{\mu4}|^2) + |U_{\mu5}|^2(1 - |U_{\mu5}|^2) - |U_{\mu4}|^2|U_{\mu5}|^2 \right]$$

$$P_{e\mu} = 2 \left[ |U_{e4}|^2|U_{\mu4}|^2 + |U_{e5}|^2|U_{\mu5}|^2 + \text{Re}(U_{e4}^* U_{\mu4} U_{e5} U_{\mu5}^*) \right]$$

It follows

$$\begin{aligned} 2P_{e\mu} &\simeq (1 - P_{ee})(1 - P_{\mu\mu}) \\ &\quad + 4 \left[ \text{Re}(U_{e4}^* U_{\mu4} U_{e5} U_{\mu5}^*) + 4|U_{e4}|^2|U_{\mu5}|^2 + 4|U_{e5}|^2|U_{\mu4}|^2 \right] \\ &= (1 - P_{ee})(1 - P_{\mu\mu}) - 2 \left[ |U_{e4}|^2|U_{\mu5}|^2 + |U_{e5}|^2|U_{\mu4}|^2 \right] \\ &\quad - 2|U_{e4} U_{\mu5} - U_{e5} U_{\mu4}|^2 \end{aligned}$$



# Relation between appearance and disappearance

## 3 + 2 neutrinos

At  $L \gg 4\pi E/\Delta m_{41}^2$ , but  $L \ll 4\pi E/\Delta m_{31}^2$

$$P_{ee} = 1 - 2 \left[ |U_{e4}|^2(1 - |U_{e4}|^2) + |U_{e5}|^2(1 - |U_{e5}|^2) - |U_{e4}|^2|U_{e5}|^2 \right]$$

$$P_{\mu\mu} = 1 - 2 \left[ |U_{\mu4}|^2(1 - |U_{\mu4}|^2) + |U_{\mu5}|^2(1 - |U_{\mu5}|^2) - |U_{\mu4}|^2|U_{\mu5}|^2 \right]$$

$$P_{e\mu} = 2 \left[ |U_{e4}|^2|U_{\mu4}|^2 + |U_{e5}|^2|U_{\mu5}|^2 + \text{Re}(U_{e4}^* U_{\mu4} U_{e5} U_{\mu5}^*) \right]$$



It follows

$$2P_{e\mu} \leq (1 - P_{ee})(1 - P_{\mu\mu})$$

Unlike in the 3 + 1 case, for 3 + 2 models, there is **NO one-to-one relation** between the appearance and disappearance probabilities.

However, there is an **inequality**, which can be used to set meaningful constraints.

# Impact of $\theta_{13}$



- Sterile neutrinos do not impact  $\theta_{13}$  measurement
- $\theta_{13} \neq 0$  does not impact sterile neutrino search

JK Machado Maltoni Schwetz, arXiv:1303.3011

# Differences between our fit and Giunti et al.


- **MiniBooNE fit**  
we use MB analysis based on official MC events, include BG oscillation
- **MINOS fit**  
we fit CC+NC data, including ND and FD, detector response matrices based on official MINOS MC
- **Reactor fit**  
minor differences in the data set, possibly different treatment of correlations among systematic uncertainties
- **LSND fit**  
Note that LSND spectral data is more constraining than the total count rate. We use this information; our fit is consistent with the numbers reported in hep-ex/0203023 (Church, Eitel, Mills, Steidl, combined LSND+KARMEN analysis)
- **Atmospheric neutrinos**  
Full fit vs. tabulated  $\chi^2$

# Hidden sector gauge forces and SBL oscillations

If sterile neutrinos have new interactions with SM fermions (e.g. in models with “baryonic sterile neutrinos”),  
new MSW potentials will influence oscillations.

How does this affect the tension in the SBL data?

Karagiorgi Shaevitz Conrad, arXiv:1202.1024  
Pospelov, arXiv:1103.3261




JK, Johannes Welter, 1408.0289

# Hidden sector gauge forces and dark matter

Interesting connection to dark matter physics:

The **same gauge force** that suppressed sterile neutrino production can also **solve small scale structure problems**:

- Too big to fail problem
- Cusp vs. core problem
- Missing satellites problem



Dasgupta JK arXiv:1310.6337

## Two further remarks

- If **sterile** and **visible** sectors have ever been in **thermal equilibrium**,  $\nu_s$  will have been **produced thermally** very early on.
- But **temperatures** of the two sectors are very different:

$$T_{\text{visible}} > T_{\text{sterile}}$$

after the SM phase transitions.

→  $\nu_s$  **abundance**  $\ll$  active neutrino abundance

## Two further remarks

- If **sterile** and **visible** sectors have ever been in **thermal equilibrium**,  $\nu_s$  will have been **produced thermally** very early on.
- But **temperatures** of the two sectors are very different:

$$T_{\text{visible}} > T_{\text{sterile}}$$

after the SM phase transitions.

→  $\nu_s$  **abundance**  $\ll$  active neutrino abundance

Mixing of  $U(1)_s$ -charged  $\nu_s$  with active neutrinos:

$$\mathcal{L} \supset -\bar{L}Y_\nu\tilde{H}\nu_R - \bar{\nu}_s Y_s H_s \nu_R - \frac{1}{2}(\nu_R)^c M_R \nu_R + h.c. ,$$

( $\tilde{H}$  = SM Higgs,  $H_s$  = sterile sector Higgs)

see e.g. Harnik JK Machado arXiv:1202.6073



# Example 1: Neutrino magnetic moments

Assume neutrinos carry an enhanced **magnetic moment**

$$\mathcal{L}_{\mu_\nu} \supset \mu_\nu \bar{\nu} \sigma^{\alpha\beta} \partial_\beta \mathbf{A}_\alpha \nu, \quad \mu_\nu \gg \mu_{\nu, \text{SM}} = 3.2 \times 10^{-19} \mu_B$$

Cross section **large** at low energies due to photon propagator  $\propto q^{-2}$

$$\frac{d\sigma_\mu(\nu e \rightarrow \nu e)}{dE_r} = \mu_\nu^2 \alpha \left( \frac{1}{E_r} - \frac{1}{E_\nu} \right),$$





# Example: A not-so-sterile 4th neutrino

Introduce a **new  $U(1)'$  gauge boson  $A'$**  (hidden photon) and a **light sterile neutrino  $\nu_s$**

Related model with gauged  $U(1)_B$  first discussed in Pospelov 1103.3261  
detailed studies in Harnik JK Machado 1202:6073 and Pospelov Pradler 1203.0545

- $\nu_s$  charged under  $U(1)'$   $\rightarrow$  direct coupling to  $A'$
- SM particles couple to  $A'$  only through kinetic mixing

$$\mathcal{L} \supset -\frac{1}{4} F'_{\mu\nu} F'^{\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} \epsilon F'_{\mu\nu} F^{\mu\nu} + \bar{\nu}_s i \not{\partial} \nu_s + g' \bar{\nu}_s \gamma^\mu \nu_s A'_\mu \\ - \overline{(\nu_L)^c} m_{\nu_L} \nu_L - \overline{(\nu_s)^c} m_{\nu_s} \nu_s - \overline{(\nu_L)^c} m_{\text{mix}} \nu_s$$

A small fraction of solar neutrinos can oscillate into  $\nu_s$

$\nu_s$  scattering cross section in the detector given by

$$\frac{d\sigma_{A'}(\nu_s e \rightarrow \nu_s e)}{dE_r} = \frac{\epsilon^2 e^2 g'^2 m_e}{4\pi p_\nu^2 (M_{A'}^2 + 2E_r m_e)^2} [2E_\nu^2 + E_r^2 - 2E_r E_\nu - E_r m_e - m_\nu^2]$$

# Temporal modulation of neutrino signals


Signals of **new light force mediators** and/or **sterile neutrinos** can show **seasonal modulation**:

- The **Earth–Sun distance**: Solar neutrino flux **peaks in winter**.

# Temporal modulation of neutrino signals

Signals of **new light force mediators** and/or **sterile neutrinos** can show **seasonal modulation**:

- The **Earth–Sun distance**: Solar neutrino flux **peaks in winter**.
- **Active–sterile neutrino oscillations**: For oscillation lengths  $\lesssim 1$  AU, sterile neutrino appearance depends on the time of year.



Harnik JK Machado, arXiv:1202.6073

# Temporal modulation of neutrino signals

Signals of **new light force mediators** and/or **sterile neutrinos** can show **seasonal modulation**:

- **Earth–Sun distance**: Solar neutrino flux **peaks in winter**.
- **Active–sterile neutrino oscillations**: For oscillation lengths  $\lesssim 1 \text{ AU}$ , sterile neutrino **appearance** depends on the time of year.
- **Sterile neutrino absorption**: For **strong  $\nu_s$ – $A'$  couplings** and **not-too-weak  $A'$ –SM couplings**, sterile neutrino **cannot traverse the Earth**.  
→ **lower flux at night**. And **nights are longer in winter**.


# Temporal modulation of neutrino signals

Signals of **new light force mediators** and/or **sterile neutrinos** can show **seasonal modulation**:

- **Earth–Sun distance**: Solar neutrino flux **peaks in winter**.
- **Active–sterile neutrino oscillations**: For oscillation lengths  $\lesssim 1 \text{ AU}$ , sterile neutrino **appearance** depends on the time of year.
- **Sterile neutrino absorption**: For strong  $\nu_s\text{--}A'$  couplings and **not-too-weak  $A'\text{--SM}$  couplings**, sterile neutrino **cannot traverse the Earth**.  
→ lower flux **at night**. And **nights are longer in winter**.
- **Earth matter effects**: An **MSW-type resonance** can lead to modified flux of certain neutrino flavors **at night**. And **nights are longer in winter**.

# Hidden photons

$$\mathcal{L} \supset -\frac{1}{4} F'_{\mu\nu} F'^{\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} \epsilon F'_{\mu\nu} F^{\mu\nu} + \bar{\nu}_s i \not{\partial} \nu_s + g' \bar{\nu}_s \gamma^\mu \nu_s A'_\mu$$



Constraints from Jaeckel Ringwald 1002.0329, Redondo 0801.1527, Biorken Essig Schuster Toro 0906.0580, Dent Ferrer Krauss 1201.2683, Harnik JK Machado

## 3+3 flavor toy model

Consider toy model with 3 sterile neutrinos, each of them mixing with only one of the active flavors:

$$U = R_{14}(\theta) R_{25}(\theta) R_{36}(\theta) U_{PMNS}, \quad R_{ij} = \text{rotation in } ij\text{-plane}.$$

Hamiltonian:

$$\mathcal{H} \simeq E + \frac{1}{2E} U \mathcal{D} U^\dagger + V_{MSW}, \quad \mathcal{D} = \text{diag}(0, \Delta m_{21}^2, \Delta m_{31}^2, \Delta m_{41}^2, \Delta m_{51}^2, \Delta m_{61}^2)$$

Mikheyev-Smirnov-Wolfenstein (MSW) potential:

$$V_{MSW} = \sqrt{2} G_F \text{diag}(n_e - n_n/2, -n_n/2, -n_n/2, 0, 0, 0),$$

$n_e$  ( $n_n$ ) = electron (neutron) number density

Oscillation probability:

$$P(\nu_\alpha \rightarrow \nu_\beta) = \left| \langle \nu_\beta | e^{-i\mathcal{H}t} | \nu_\alpha \rangle \right|^2$$